

NMR structure of the N-terminal domain of the replication initiator protein DnaA

Thomas J. Lowery · Jeffrey G. Pelton · John-Marc Chandonia ·
Rosalind Kim · Hisao Yokota · David E. Wemmer

Received: 7 March 2007 / Accepted: 17 July 2007
© Springer Science+Business Media B.V. 2007

Abstract DnaA is an essential component in the initiation of bacterial chromosomal replication. DnaA binds to a series of 9 base pair repeats leading to oligomerization, recruitment of the DnaBC helicase, and the assembly of the replication fork machinery. The structure of the N-terminal domain (residues 1–100) of DnaA from *Mycoplasma genitalium* was determined by NMR spectroscopy. The backbone r.m.s.d. for the first 86 residues was 0.6 ± 0.2 Å based on 742 NOE, 50 hydrogen bond, 46 backbone angle, and 88 residual dipolar coupling restraints. Ultracentrifugation studies revealed that the domain is monomeric in solution. Features on the protein surface include a hydrophobic cleft flanked by several negative residues on one side, and positive residues on the other. A negatively charged ridge is present on the opposite face of the protein. These surfaces may be important sites of interaction with other proteins involved in the replication process. Together, the structure and NMR assignments should facilitate the design of new experiments to probe the protein–protein interactions essential for the initiation of DNA replication.

Keywords Replication · DnaB · DiaA · ATPase · Initiation

Thomas J. Lowery and Jeffrey G. Pelton contributed equally to this work.

T. J. Lowery · J. G. Pelton · J.-M. Chandonia · R. Kim ·
H. Yokota · D. E. Wemmer
Lawrence Berkeley National Laboratory, 1 Cyclotron Rd,
Berkeley, CA 94720, USA

T. J. Lowery · D. E. Wemmer (✉)
Department of Chemistry, University of California at Berkeley,
Berkeley, CA 94720, USA
e-mail: DEWemmer@lbl.gov

The bacterial protein DnaA plays a central role in the initiation of chromosomal replication and is involved in the activation and repression of several genes, including its own gene (*dnaA*) in *E. coli*. Replication begins with the formation of oligomers of DnaA on recognition sites at the origin of replication (OriC). The DnaA complex unwinds and opens several AT-rich segments of double-stranded DNA within the origin and stabilizes the resulting single-strands. DnaA then recruits the DnaB helicase with help from DnaC to form the “prepriming complex” [1]. After priming of the single-stranded DNA, replication occurs by the action of DNA Polymerase III. ATP hydrolysis acts as the switch between active (ATP-bound) and inactive (ADP-bound) forms of DnaA. Initiation begins only with the ATP-bound form (for reviews see [2–4]).

The DnaA protein is composed of four functionally independent domains [4, 5]. Domains III and IV exhibit ATPase and DNA-binding activities, respectively, and are more highly conserved than the N-terminal domains I and II [3]. Evidence suggests that domains I and III are involved in oligomerization and binding DnaB, and that domain II is a flexible linker. Recent X-ray structures of domains III and IV of DnaA from *Aquifex aeolicus* bound to ADP [6] and a nonhydrolyzable ATP-analog [7], as well as domain IV bound to DNA [8] showed that domain III resembles other members of the AAA+ family of ATPases, and that domain IV adopts a helix-turn-helix DNA-binding fold. In contrast, the secondary structure of domain I has been predicted based on multiple sequence alignments [5, 9, 10], however, little is known about its tertiary structure.

DnaA recognizes five high affinity 9 base pair sites as well as several lower affinity 6 base pair sites within OriC via the C-terminal helix-turn-helix DNA-binding domain (domain IV) [11, 12]. In addition to direct binding, the formation of oligomers containing 20–30 molecules of

DnaA at the origin has been demonstrated by electron microscopy [13, 14]. The mechanism of oligomer formation has been studied extensively [3, 15], but the nature of the complex remains poorly understood. Replacement of the dimerization domain of the lambda cI repressor with the N-terminal domain of DnaA resulted in a functional chimeric protein, suggesting direct interactions between the N-terminal domains [10, 16]. Results from solid phase binding assays [10] and gel mobility shift experiments [17] have also suggested interactions between N-terminal domains. In the latter study, the N-terminal domain was shown to be required for cooperative binding of the protein to adjacent DnaA boxes. Mutational analysis has implicated Leu5, Trp6, Glu8, and Cys9 in domain I of *E. coli* DnaA as being critical for self-oligomerization [18, 19]. Other experiments have suggested a role for the ATPase domain in oligomerization [6, 7, 15, 16, 20]. Together, the observations that full-length DnaA from *E. coli* exists as a monomer in both the ADP- and ATP-bound forms, as well as in the presence of DNA that contains a single high affinity binding site [9, 12], have made it difficult to characterize the exact nature of self-oligomerization.

In addition to oligomerization, cross-linking studies have shown that the N-terminal domain of DnaA interacts directly with both DnaB [1] and the regulatory protein DiaA [21]. For DnaB, the regions of interaction were further defined by solid phase binding assays, which showed that residues 24–86 from the N-terminal domain, as well as residues in domain III, were important [20]. Within this set of residues, Pro28 appears to be especially important [19].

As part of a structural genomics effort we developed an algorithm to identify folded domains within larger proteins that had no known structural homologs [22]. The sequence corresponding to the N-terminal segment (residues 1–100) of DnaA from *Mycoplasma genitalium* was one such domain candidate. Initial NMR results indicated that this construct was indeed folded and a good candidate for structure determination. Herein, we present the structure of this N-terminal domain of DnaA, show that it exists as a monomer in solution, and relate it to previous genetic and biochemical data. The results presented provide a structural framework needed for the design of new experiments to further our understanding of the initiation of DNA replication.

Materials and methods

The sequence containing domain I and a portion of domain II of DnaA from *M. genitalium* (Swiss-Prot entry DNAA_MYCGE, accession code P35888; PFAM id PF003408; TIGR id MG469, gi number 12045329; TIGR-FAM id TIGR00362; InterPro id IPR001957; residues 1–

100) was selected and cloned as described [22, 23]. The protein was expressed as an N-terminal 6-His-TEV fusion protein in *E. coli* BL21* (DE3). In this construct an N-terminal methionine and six glycine residues precede the first methionine of DnaA. These residues were not assigned or included in the structure calculations. Unlabeled protein was obtained by growing the cells in 1 l of LB at 37°C to an OD₆₀₀ of 0.6 followed by induction with 1 mM isopropylthio-β-D-galactoside (IPTG) for 3.5 h. The molecular weight of the resulting protein was within one mass unit of the expected weight. Labeled protein was obtained by growing cells in 1 l of LB to an OD of 0.7, followed by pelleting the cells and transferring them to 0.75 l of isotope labeled M9 as described [24]. Isotopically labeled M9 contained either ¹⁵N-labeled ammonium chloride or both ¹⁵N-labeled ammonium chloride and ¹³C-labeled glucose, to produce ¹⁵N- and ¹⁵N/¹³C-labeled proteins, respectively. The cells were lysed in buffer containing 50 mM sodium phosphate (pH 7.5), 0.1 M sodium chloride, 1 mM PMSF, protease inhibitor cocktail Roche Molecular Biochemicals, Mannheim, Germany), and cell debris was pelleted by ultracentrifugation. The soluble His-tagged protein was purified by nickel affinity chromatography using a running buffer of 50 mM sodium phosphate (pH 7.5), 0.1 M sodium chloride, 5% v/v glycerol, and 5 mM imidazole. After several column washes, the protein was eluted by increasing the imidazole concentration from 5 mM to 1 M. After concentration to approximately 2 ml by ultracentrifugation (5 kD MWCO Amicon Ultra), further purification was performed using size exclusion chromatography (HiLoad 16/60 Superdex 75pg; GE Healthcare) with a running buffer containing 50 mM sodium phosphate (pH 7.5) and 0.15 M sodium chloride. Fractions containing the N-terminal domain were pooled and treated with TEV protease overnight at room temperature. The protein was then reapplied to a nickel-affinity column and the eluate was collected and concentrated. Analytical size exclusion chromatography was conducted on a BioCad Sprint II with a TSK-Gel G4000SW_{xl} column (Tosoh Bioscience LLC; 7.8 mm ID × 30 cm) using bovine serum albumin, carbonic anhydrase, and lysozyme as molecular weight standards. Analytical ultracentrifugation (AUC) was performed using a Beckman Optima XL-I analytical ultracentrifuge. Purified protein was dialyzed against 50 mM sodium phosphate (pH 6.8) and 0.1 M sodium chloride overnight at 4°C. Protein at a concentration of 140 μM was loaded into each of five sample cells (100 μl) as determined by UV absorbance ($\epsilon_{280} = 5.12 \text{ mM}^{-1} \text{ cm}^{-1}$). Blank cells were filled with dialysis buffer (110 μl). The centrifuge was allowed to equilibrate for 15 h at 30,000 rpm and 20°C before the recording of 10 sequential scans. Data were fit using UltraScan software [25]. NMR samples were prepared by concentrating the purified protein in buffer containing

50 mM sodium phosphate (pH 6.8), 0.1 M sodium chloride, 2 mM EDTA, 0.05% azide, protease inhibitor cocktail (Roche Molecular Biochemicals, Mannheim, Germany), and 0.1 mM PMSF to a concentration of 1.0–1.5 mM. NMR samples in D₂O were prepared by lyophilizing an existing NMR sample and re-dissolving it into D₂O. NMR spectra were recorded on Bruker NMR spectrometers, including a DRX 500 equipped with a cryoprobe, a DMX 600, an Avance 800 equipped with a cryoprobe, and an Avance II 900. All spectra were recorded at 25°C. Backbone resonances were assigned with 3D HNCA [26], HNCOC [27, 28], CBCA(CO)NH [29], HNCACB [30] spectra recorded at 500 MHz and HBHACONH [31] and HNCACO [27] spectra recorded at 600 MHz. Side-chain aliphatic ¹³C/¹H signals were assigned with 3D-HBHA(-CO)NH, 3D-(H)C(CO)NH [32], 3D-H(CCO)NH [32], 3D-¹⁵N TOCSY [33], and 3D-HCCH-TOCSY [34] spectra at 600 MHz. In the final analysis, all of the backbone ¹H, ¹⁵N, ¹³C α , and ¹H α signals and 88% of the sidechain signals were assigned, excluding the N-terminal methionine and six glycine residues that result from cloning, as well as the first methionine residue (Met1). Phi torsion angle restraints were obtained from a 3D HNHA spectrum [35] recorded at 800 MHz. Stereospecific assignments for the methyl groups of 3 of 4 valine and 11 of 12 leucine residues were obtained by comparison of HSQC spectra of fully and 10% ¹³C-labeled samples [36, 37]. NOEs were identified in 3D ¹⁵N-¹H NOESY-HSQC (800 MHz), 3D ¹³C-¹H NOESY-HSQC (800 and 900 MHz), and a 4D ¹³C-HMQC-NOESY-HSQC [38] (600 MHz) spectra with mixing times ranging between 85 and 100 ms. A total of 53 ¹H-¹⁵N and 35 ¹H-¹³C α residual dipolar couplings (rdcs) were used in the structure calculations. The ¹H-¹⁵N rdcs were obtained from IPAP [39] spectra recorded on a uniformly ¹⁵N-labeled protein sample, and the ¹H-¹³C α dipolar couplings were based on a 3D (HA)CA(CO)NH spectrum [40], recorded on a uniformly-¹⁵N/¹³C labeled protein sample. Both proteins were dissolved in 12 mg/ml pf1 phage [41]. Tensor parameters were determined from histogram plots of the couplings [42] and values based on intermediate structures [43, 44]. The magnitude of the alignment tensor and rhombicity were set to -13.7 Hz and 0.47 for HN vectors and -13.3 Hz and 0.47 for C α H vectors. Force constants were adjusted to 0.4 kcal mol⁻¹ Hz⁻² and 0.2 kcal mol⁻¹ Hz⁻² to yield r.m.s.d.s equal to the uncertainties in the measurements (Table 1). Structures were calculated using a torsion-angle simulated annealing protocol [51, 52] as implemented in the XPLOR-NIH program, version 2.11.2 [53]. Center averaging and distance corrections [54] were used for non-stereoassigned protons. The NOE and rdc data were generally consistent with one another as evidenced by the fact that average structures (20 best of 200) calculated with and without rdcs superimposed with an

r.m.s.d of 0.67 Å. The main difference was a slight displacement of the end of helix III at Asn53. The number of restraints and structure statistics are presented in Table 1.

Results and discussion

As discussed above, the results from several studies suggested that the N-terminal domain dimerizes or assists in the formation of larger oligomers [10, 18, 19]. Before initiating NMR studies, we probed the oligomerization state of our construct using both size exclusion chromatography (SEC) and analytical ultracentrifugation (AUC). The initial SEC results suggested that the protein was a dimer at a concentration of 1 mM (data not shown). In contrast, the AUC results, which were conducted at a concentration of 140 μ M, revealed that the protein was monomeric (data not shown). In addition, identical ¹H-¹⁵N HSQC spectra were obtained at 70 μ M and the working concentration of 1.2 mM (Fig. 1), indicating that protein exists in the same state at both concentrations. Because results from AUC depend only on the mass of the protein, whereas results from SEC are influenced by the shape of the molecule, we conclude that the protein is a monomer in solution at millimolar concentrations, and that the apparent molecular weight determined from SEC was an artifact, likely resulting from the first 6 and last 15 residues being unstructured and flexible in solution (see below).

A summary of the NMR-derived restraints and structural statistics are presented in Table 1. The NMR data reveal that domain I is composed of five helices and a three-stranded β -sheet with topology $\alpha\alpha\beta\alpha\alpha\beta\alpha$ (Fig. 2). The experimentally determined secondary structure is in general agreement with predictions based on multiple sequence alignments [5, 9, 10]. The exceptions are the presence of β -strand 1 (residues 31–34), a break between helices III and IV, and the presence of helix V (residues 77–84). Helix V may not be present in *E. coli* since there are two proline residues (Pro82 and Pro85) in this region. The three-dimensional structure is formed by the packing of helices I and IV and II and III against one another and one side of the β -sheet, while helix V packs against the opposite face (Fig. 3). As mentioned previously, the last fifteen residues of the construct, which correspond to the first residues of domain II [4], show no long-ranges NOEs, suggesting that they are unstructured. These data are consistent with this segment functioning as a flexible linker between domain I and the ATPase and HTH domains [4]. A search for structural homologs [55] revealed that the N-terminal domain is similar to ribosomal binding factor A (RbfA) [56, 57], which contains a type II KH RNA-binding fold (Z score of 3–4). However, the structural similarity

Table 1 Structural statistics for the N-terminal domain of DnaA from *M. genitalium*^a

RMSD from restraints ^b			
NOE (Å)			0.014 ± 0.002
Cdih (°)			0.37 ± 0.09
HN rdc R-factor (%) /rmsd (Hz)			2.5 ± 0.3 / 0.46 ± 0.06
C ^α H rdc R-factor (%) /rmsd (Hz)			9.5 ± 0.7% / 1.7 ± 0.1
RMSD from idealized geometry			
Bond (Å)			0.00174 ± 0.00007
Angle (°)			0.426 ± 0.006
Improper (°)			0.25 ± 0.008
Number of violations/structure			
NOE (0.1 Å–0.3 Å)			1.9 ± 1.5
NOE (>0.3 Å)			0
Cdih (>5°)			0
HN rdc (>1.5 Hz)			0.3 ± 0.07
C ^α H RDCs (>3 Hz)			3 ± 1
Coordinate RMSDs to the mean ^c			
Secondary structure/backbone (Å)			0.6 ± 0.2
Residues 3–86/backbone (Å)			0.7 ± 0.2
Heavy atom < 40% solvent exposed (Å)			1.2 ± 0.1
Structure quality factors—overall statistics ^d			
	Mean score (all models)	Standard Deviation	Z-score ^e
Procheck G-factor (phi-psi only) ^f	−1.04	N/A	−3.78
Procheck G-factor (all dihedral) ^f	−1.06	N/A	−6.27
Verify3D	0.25	0.03	−3.37
Prosall (-ve)	0.34	0.05	−1.28
MolProbity clashscore	58	6	−8.47
Ramachandran Plot Summary for ordered residues ^{d,f}			
Most favored regions (%)			70.7
Additionally allowed regions (%)			28.6
Generously allowed regions (%)			0.7
Disallowed regions (%)			0.0

^a Parameters are for the 20 best of 200 structures

^b A total of 286 intra-residue, 194 sequential, 103 medium-range, and 159 long-range NOEs, 50 hydrogen bonds (2 per hydrogen bond), 46 phi angle restraints, 53 HN residual dipolar couplings (rdcs), and 35 C^αH rdc were used in the calculations. The RDC R-factor is defined as the ratio of the r.m.s. deviation between observed and calculated values to the expected r.m.s. deviation if the vectors were randomly distributed [45]

^c Secondary structure included N,C^α,C' atoms for helical regions 3–14, 19–24, 46–52, 59–62, 77–85, and β-strand regions 32–34, 39–42, 72–75. Less than 40% solvent exposed residues included 4, 6–11, 14–16, 18–23, 25–29, 31–34, 36–43, 45, 48, 49, 51–53, 55–59, 61–66, 68, 69, 71–77, 79–81, 83–85

^d Determined using Protein Structure Validation Software version 1.3 [46]. Software included in the calculations includes Procheck [47], Verify3D [48], Prosall [49], and MolProbity [50]

^e With respect to mean and standard deviation for a set of 252 X-ray structures of less than 500 residues, of resolution less than or equal to 1.80 Å, R-factor less than or equal to 0.25 and R-free less than or equal to 0.28; a positive value indicates a 'better' score

^f Residues 2–26, 30–33, 38–55, 58–63, 69–85

probably does not carry over to function. To our knowledge, there is no evidence that the N-terminal domain of DnaA binds to nucleic acids.

A total of 17 residues are conserved among the roughly 212 known sequences of the N-terminal domain of DnaA (Fig. 2). Approximately 80% of these residues are involved in hydrophobic packing within the protein core (Fig. 3b).

Three other conserved residues, Asn45, Asn53, and Glu57, are located on the surface of the protein in helices III and IV. Comparison also shows that proline residues in *E. coli* and the consensus sequence (not shown) map to loops and unstructured regions of the structure (Fig. 2). The only exceptions are the two proline residues in *E. coli* that map to helix V, as mentioned above. Similarly, the locations of

Fig. 1 ^1H - ^{15}N HSQC spectra recorded at 500 MHz of the N-terminal domain of DnaA from *M. genitalium* at concentrations of (a) 70 μM and (b) 1.2 mM. Chemical shift assignments are denoted by one letter amino acid code and residue number in (b). The peak labeled with U is unassigned. The position of the peak is consistent with one or more of the 6 glycine residues at the N-terminus that result from a cloning artifact





Fig. 2 Sequences of the N-terminal domains of DnaA from *M. genitalium* and *E. coli*. The alignment is based on 212 DnaA sequences. The domain boundaries are taken from Messer et al. 1999 [4]. Domain III begins at residue 134 in *E. coli*. Helices and β -strands found in DnaA from *M. genitalium* are shown schematically above the sequences as black bars and arrows, respectively. Asterisks (*)

denote the positions of conserved residues within the multiple sequence alignment, carets (^) denote the locations of proline residues in *E. coli* or the consensus sequence, and bars (|) indicate the positions of insertions in other DnaA sequences. Hydrophobic residues common in both *M. genitalium* and *E. coli* are shaded grey

Fig. 3 (a) Backbone trace for the top 20 structures superimposed using the backbone N, C α , and C' coordinates. Unstructured residues (89–100) are not shown. (b) Ribbon diagram of the structure. The locations of conserved hydrophobic residues based on the full multiple sequence alignment are shown in yellow. Sites of functionally critical residues are denoted with black arrows. (c) Electrostatic surface of the structure showing positively charged regions in blue, neutral regions in white, and negatively charged regions in red. (c—left) The hydrophobic cleft near helix I is apparent, with positive residues along the bottom edge and negative residues along the top edge (see text). (c—right) A 180 degree rotation of the molecule shows the negatively charged surface associated with helices III and IV. Figure 3b, c were made with PyMOL (Delano Scientific, Palo Alto, Ca)



insertions in other sequences map to loops or unstructured regions. For insertions, the only exceptions occur in three mycobacterium sequences, in which there is an insertion in helix II. Together, these observations indicate that the overall fold of the N-terminal domain is conserved across all species.

As was previously noted [5, 9, 10], the sequence similarity within domain I is lower than within domains III and IV, making it difficult to identify functionally important residues. Experimentally, residues at positions Gln3, Phe4, and Phe7 (Leu5, Trp6 and Cys9 in *E.coli*) have been shown to be critical for oligomerization [18, 19] and residues 24–86, and in particular Pro28, have been implicated in DnaB binding [19, 20]. Residues Gln3, Phe4, and Phe7 line the inner face of helix I (Fig. 3) and Pro28 is in the loop connecting helix II and β -strand I. These secondary structure elements form a hydrophobic cleft with positive residues along the bottom edge (Gln3, Lys8, Lys12, and Lys13), and negative residues along the top edge (Asp23 and Asp28) (Fig. 3c). Approximately half of the residues lining this cleft are conserved in the full multiple sequence alignment. On the opposite face of the protein, a negatively charged ridge runs from Glu46 to Glu67 along the axes of helices III and IV (Fig. 3c). These surface features may be

important in the formation of DnaA oligomers and binding with DnaB, DiaA, and other proteins involved in the replication process.

DnaA oligomerizes at the origin of replication and interacts with a host of other proteins, including DnaB and DiaA, during DNA replication [2, 3]. To date, the nature of these interactions and the specific role of the N-terminal domain remain poorly understood. The chemical shift assignments presented here should promote efforts to explore these interactions directly at the molecular level. In addition, the structure and surface analysis should enable the design of more precise genetic studies, which, together with binding studies, should help unravel the complex set of protein–protein interactions that are central to this fundamental process.

Assession codes

NMR restraints and coordinates have been deposited in the Protein Data Bank under accession number 2JMP. Chemical shift assignments have been deposited in the BioMagResBank (BMRB) under accession number BMRB-15055.

Addendum

After submission of this manuscript, the NMR structure of the N-terminal domain of DnaA from *E. coli* was reported [58]. The overall fold is similar to the structure reported here, with a pairwise coordinate RMSD of 2.9 Å over 76 residues (14% sequence identity), obtained using the Dali algorithm [59]. The biggest difference is the lack of the C-terminal helix in the *E. coli* structure compared to that from *M. genitalium*, as predicted based on sequence alignment [5, 9, 10].

Acknowledgements We thank D. King for mass spectral analysis, and J. Erzberger, M. Mott, and J. Berger for helpful discussions. We also thank M. Doucleff for recording several spectra. T.J.L. acknowledges the University of California Biotechnology Research and Education Program for financial support. This work was supported by grant GM62412 to the Berkeley Structural Genomics Center from the National Institute of General Medical Sciences, National Institutes of Health. We also thank NSF (BBS87W134 and 0119304) and NIH (RR15756 and GM068933) for funding NMR equipment.

References

- Marszalek J, Kaguni JM (1994) *J Biol Chem* 269:2883–2890
- Kato J (2005) *Crit Rev Biochem Mol Biol* 40:331–342
- Messer W (2002) *FEMS Microbiol Rev* 26:355–374
- Messer W, Blaesing F, Majika J, Nardmann J, Schaper S, Schmidt A, Seitz H, Speck C, Tuengler D, Wegrzyn G, Weigel C, Welzcek M, Zakrzewska-Czerwinska J (1999) *Biochimie* 81:819–825
- Sutton MD, Kaguni JM (1997) *J Mol Biol* 274:546–561
- Erzberger JP, Pirruccello MM, Berger JM (2002) *EMBO J* 21:4763–4773
- Erzberger JP, Mott MI, Berger JM (2006) *Nat Struct Biol* 13:676–683
- Fujikawa N, Kurumizaka H, Nureki O, Terada T, Shirouzu M, Katayama T, Yokoyama S (2003) *Nucleic Acids Res* 31:2077–2086
- Schaper S, Messer W (1995) *Proteins* 28:1–9
- Weigel C, Schmidt A, Seitz H, Tuengler D, Welzcek M, Messer W (1999) *Mol Microbiol* 34:53–66
- Speck C, Messer W (2001) *EMBO J* 20:1469–1476
- Speck C, Weigel C, Messer W (1999) *EMBO J* 18:6169–6176
- Crooke E, Thresher R, Hwang DS, Griffin J, Kornberg A (1993) *J Mol Biol* 233:16–24
- Funnell BE, Baker TA, Kornberg A (1987) *J Biol Chem* 262:10327–10334
- Messer W, Blaesing F, Jakimowicz D, Krause M, Majka J, Nardmann J, Schaper S, Seitz H, Speck C, Weigel C, Wegrzyn G, Welzcek M, Zakrzewska-Czerwinska J (2001) *Biochimie* 83:5–12
- Jakimowicz D, Majka J, Konopu G, Wegrzyn G, Messer W, Schrepf H, Zakrzewska-Czerwinska J (2000) *J Mol Biol* 298:351–364
- Majka J, Zakrzewska-Czerwinska J, Messer W (2001) *J Biol Chem* 276:6243–6252
- Felczak M, Simmons LA, Kaguni JM (2005) *J Biol Chem* 280:24627–24633
- Simmons LA, Felczak M, Kaguni JM (2003) *Mol Microbiol* 49:849–858
- Seitz H, Weigel C, Messer W (2000) *Mol Microbiol* 37:1270–1279
- Ishida T, Akimitsu N, Kashioka T, Hatano M, Kubota T, Ogata Y, Sekimizu K, Katayama T (2004) *J Biol Chem* 279:45546–45555
- Chandonia JM, Kim SH, Brenner SE (2006) *Proteins* 62:356–370
- Shin DH, Kim JS, Yokota H, Kim R, Kim SH (2006) *Protein Sci* 15:921–928
- Marley J, Lu M, Bracken C (2001) *J Biomol NMR* 20:71–75
- Demeler G (2005). In: Scott DJ, Rowe AJ (eds) *Modern analytical ultracentrifugation: techniques and methods*, Royal Society of Chemistry, UK, pp 210–229
- Grzesiek S, Bax A (1992) *J Magn Reson* 99:201–207
- Kay LE, Xu GY, Yamazaki T (1994) *J Magn Reson* 109 A:129–133
- Stonehouse J, Clowes RT, Shaw GL, Keeler J, Laue ED (1995) *J Biomol NMR* 5:226–232
- Grzesiek S, Bax A (1992) *J Am Chem Soc* 114:6291–6293
- Wittekind M, Mueller L (1993) *J Magn Reson* 101 B:201–205
- Grzesiek S, Bax A (1993) *J Biomol NMR* 3:185–204
- Grzesiek S, Anglister J, Bax A (1993) *J Magn Reson* 101 B:114–119
- Driscoll PC, Clore GM, Marion D, Wingfield PT, Gronenborn AM (1990) *Biochemistry* 29:3542–3556
- Kay LE, Xu GY, Singer A, Muhandram D, Forman-Kay J (1993) *J Magn Reson* 101 B:333–337
- Kuboniwa H, Grzesiek S, Delaglio F, Bax A (1994) *J Biomol NMR* 4:871–878
- Neri D, Szyperski T, Otting G, Senn H, Wuthrich K (1989) *Biochemistry* 28:7510–7516
- Szyperski T, Neri D, Leiting B, Otting G, Wuthrich K (1992) *J Biomol NMR* 2:323–334
- Vuister GW, Clore GM, Gronenborn AM, Powers R, Garrett DS, Tschudin R, Bax A (1993) *J Magn Reson* 101 B:210–213
- Ottiger M, Delaglio F, Bax A (1998) *J Magn Reson* 131:373–378
- Tjandra N, Bax A (1997) *J Am Chem Soc* 119:9576–9577
- Hansen MR, Mueller L, Pardi A (1998) *Nat Struct Biol* 5:1065–1074
- Clore GM, Gronenborn AM, Bax A (1998) *J Magn Reson* 133:216–221
- Losonczi JA, Andrec M, Fischer MW, Prestegard JH (1999) *J Magn Reson* 138:334–342
- Zweckstetter M, Bax A (2000) *J Am Chem Soc* 122:3791–3792
- Bax A, Kontaxis G, Tjandra N (2001) *Methods Enzymol* 339:127–174
- Bhattacharya A, Tejero R, Montelione GT (2007) *Proteins* 66:778–795
- Laskowski RA, Rullmann JA, MacArthur MW, Kaptein R, Thornton JM (1996) *J Biomol NMR* 8:477–486
- Luthy R, Bowie JU, Eisenberg D (1992) *Nature* 356:83–85
- Sippl MJ (1993) *Proteins* 17:355–362
- Lovell SC, Davis IW, Arendall WB 3rd, de Bakker PI, Word JM, Prisant MG, Richardson JS, Richardson DC (2003) *Proteins* 50:437–450
- Rice LM, Brunger AT (1994) *Proteins* 19:277–290
- Stein EG, Rice LM, Brunger AT (1997) *J Magn Reson* 124:154–164
- Schwieters CD, Kuszewski JJ, Tjandra N, Clore GM (2003) *J Magn Reson* 160:65–73
- Fletcher CM, Jones DNM, Diamond R, Neuhaus D (1996) *J Biomol NMR* 8:292–310
- Holm L, Sander C (1993) *J Mol Biol* 233:123–138
- Huang YJ, Swapna GV, Rajan PK, Ke H, Xia B, Shukla K, Inouye M, Montelione GT (2003) *J Mol Biol* 327:521–536
- Rubin SM, Pelton JG, Yokota H, Kim R, Wemmer DE (2003) *J Struct Funct Genomics* 4:235–243
- Abe Y, Jo T, Matsuda Y, Matsunaga C, Katayama T, Ueda T (2007) *J Biol Chem* 282:17816–17827
- Holm L, Park J (2000) *Bioinformatics* 16:566–567